

CONE-FIELDS, DOMINATION, AND HYPERBOLICITY

SHELDON NEWHOUSE

1. INTRODUCTION

We obtain new conditions for dominated and hyperbolic splittings on compact invariant sets for a diffeomorphism in terms of its induced action on a cone field and its complement. The results are applied to give simple new proofs of well-known theorems for hyperbolicity of the set of bounded orbits in real and complex Henon mappings. In this latter case, much more complete information can be given making use of complex methods. See, for instance, [1]

Let M be a compact C^∞ Riemannian manifold, let $f \in \mathcal{D}^1(M)$, let Λ be a compact f -invariant set.

The notion of a dominated splitting arose in the work of Mane on the stability conjecture [4] and has been considered in several recent works. See for instance [9].

The usual definition is the following.

Definition. A *dominated splitting* on Λ is a splitting $T_x M = E_{1x} \oplus E_{2x}$ for each $x \in \Lambda$ such that there are constants $C > 0$ and $\lambda > 1$ such that

$$(1) \quad |Df_x^n|_{E_{2x}} || Df_{f^n x}^{-n}|_{E_{1f^n x}}| \leq C\lambda^{-n} \text{ for } n \geq 0,$$

$$(2) \quad Df(E_{ix}) = E_{i,fx} \text{ for } i = 1, 2, \text{ and}$$

$$(3) \quad \text{the maps } x \rightarrow E_{ix} \text{ are continuous.}$$

Let us give an interpretation of these estimates.

For a linear map $L : \mathbf{R}^n \rightarrow \mathbf{R}^n$, and a subset $E \subset \mathbf{R}^n$ with $E \neq \{0\}$, set

$$|L|E| = \sup_{v \neq 0, v \in E} \frac{|Lv|}{|v|},$$

and

$$m(L|E) = \inf_{v \neq 0, v \in E} \frac{|Lv|}{|v|}.$$

If L is an isomorphism, and E is a subspace, then

$$m(L|E) = |L^{-1}|L(E)|^{-1}.$$

We call $m(L|E)$ the *minimal expansion* of L on E , and $m(L^{-1}|L(E))$ the *minimal inverse expansion* of L on E .

We may write condition (1) in several equivalent ways. For $n \geq 0$,

$$(4) \quad |Df_x^n | E_{2x} | m(Df_x^n | E_{1x})^{-1} \leq C\lambda^{-n}$$

$$(5) \quad |Df_x^n | E_{2x} | \leq C\lambda^{-n} m(Df_x^n | E_{1x})$$

$$(6) \quad m(Df_{f^n x}^{-n} | E_{2, f^n x})^{-1} m(Df_x^n | E_{1x})^{-1} \leq C\lambda^{-n}$$

$$(7) \quad m(Df_x^n | E_{1x}) m(Df_{f^n x}^{-n} | E_{2, f^n x}) \geq C^{-1} \lambda^n.$$

The last formulation (7) can be simply expressed as follows: the minimal expansion of Df_x^n on E_{1x} times the minimal inverse expansion of Df_x^n on E_{2x} grows exponentially with n .

Remark:

- (1) Dominated splittings need not be unique. For example, consider a toral automorphism with a center and expanding subspace, or, more generally, any partially hyperbolic set.
- (2) We will show later that conditions (1) and (2) imply condition (3).
- (3) If $\dim E_{1x}$ is constant, then it follows from Theorem 1.2 below that the dominated splitting is unique in the following sense. If $T_x = F_{1x} \oplus F_{2x}$ is another splitting of $T_x M$ such that, for all $x \in \Lambda$,

$$m(Df_x^n | F_{1x}) m(Df_{f^n x}^{-n} | F_{2, f^n x}) \geq C^{-1} \lambda^n$$

and

$$\dim E_{1x} = \dim F_{1x},$$

then $E_{1x} = F_{1x}$ and $E_{2x} = F_{2x}$ for all $x \in \Lambda$.

In this note we are interested in establishing simple conditions which guarantee that dominated splittings exist. Our conditions will be in terms of the action of Df on certain cone fields over the set Λ . In many examples, these conditions are easy to verify. They also lead to an elegant and useful method for proving that invariant sets are uniformly hyperbolic. This has many applications in bifurcation theory and other areas. As a particular application, we give a simple new proof of hyperbolicity of the set of bounded orbits for certain Henon mappings in both the real and complex situations.

There are many known results containing sufficient conditions for hyperbolicity. See, for instance, the papers [3], [7], and the books [6], [8], and [10]. Our techniques here are analogous to some which appear in [8] and [10] for two dimensional systems involving invariant cone fields. We replace the invariance conditions in [8] and [10] by expansion conditions on the cones and their complements. This gives stronger results and fits in easily in the more general context of domination. The techniques in this section are simple, and, although we expect that they are known to experts, we have found no reference with analogous results in the generality presented here.

Let us begin by recasting the domination conditions in a way which gives uniqueness and continuity.

Let $E \subset \mathbf{R}^n$ be a proper subspace; i.e., $0 < \dim E < n$. Let F be a complementary subspace; i.e., $\mathbf{R}^n = E \oplus F$.

The *standard unit cone* determined by the subspaces E and F is the set

$$K_1(E, F) = \{v = (v_1, v_2) : v_1 \in E, v_2 \in F, \text{ and } |v_2| \leq |v_1|\}.$$

A *cone* in \mathbf{R}^n with core E , denoted $\mathcal{C}(E)$, is the image $T(K_1(E, F))$ where $T : \mathbf{R}^n \rightarrow \mathbf{R}^n$ is a linear automorphism such that $T(E) = E$. By a **cone** \mathcal{C} in \mathbf{R}^n we mean a set $\mathcal{C}(E)$ for some proper subspace E of \mathbf{R}^n .

Now consider a diffeomorphism f on the manifold M and compact invariant set Λ as above.

A *cone field* $\mathcal{C} = \{\mathcal{C}_x\}$ on Λ is a collection of cones $\mathcal{C}_x \subset T_x M$ for $x \in \Lambda$. We say that the cone field \mathcal{C}_x has *constant orbit core dimension* on Λ if $\dim E_x = \dim E_{f_x}$ for all $x \in \Lambda$ where E_x, E_{f_x} are the cores of $\mathcal{C}_x, \mathcal{C}_{f_x}$, respectively. Note that there is no a priori continuity assumption on $x \rightarrow \mathcal{C}_x$.

Given such a cone field $\mathcal{C} = \{\mathcal{C}_x\}_{x \in \Lambda}$, let

$$m_{\mathcal{C},x} = m_{\mathcal{C},x}(f) = \inf_{v \in \mathcal{C}_x \setminus \{0\}} \frac{|Df_x(v)|}{|v|},$$

and

$$m'_{\mathcal{C},x} = m'_{\mathcal{C},x}(f) = \inf_{v \notin \mathcal{C}_{f_x}} \frac{|Df_{f_x}^{-1}(v)|}{|v|}.$$

We call $m_{\mathcal{C},x}$ the *minimal expansion* of f on \mathcal{C}_x or of Df on \mathcal{C}_x , and we call $m'_{\mathcal{C},x}$ the *minimal co-expansion* of f on \mathcal{C}_x or of Df on \mathcal{C}_x .

We define the *domination coefficient* of f on \mathcal{C} to be

$$(8) \quad m_d(\mathcal{C}) = m_d(\mathcal{C}, f) = \inf_{x \in \Lambda} m_{\mathcal{C},x} \cdot m'_{\mathcal{C},x}.$$

We say that f is *dominating* on \mathcal{C} over Λ (or \mathcal{C} is a *dominating cone field* on Λ) if \mathcal{C} has constant orbit core dimension and $m_d(\mathcal{C}) > 1$.

Remark. This condition depends on the choice of Riemannian metric on M . An analogous weaker condition, which is independent of the Riemannian metric, is that $m_d(\mathcal{C}, f^{n_0}) > 1$ for some $n_0 \geq 1$. There are obvious modifications of the results below which hold under this weaker condition.

We have the following simple lemma.

Lemma 1.1. *Suppose that f is dominating on \mathcal{C} over Λ . Then, \mathcal{C} is an f -invariant cone field. That is, for $x \in \Lambda$, we have*

$$(9) \quad Df_x(\mathcal{C}_x) \subset \mathcal{C}_{f_x}.$$

Proof.

Let $v \in \mathcal{C}_x \setminus \{0\}$, and let $w = Df_x(v)$.

Then,

$$|Df_x(v)| \geq m_{\mathcal{C},x}(f) |v|.$$

We want to show that $w \in \mathcal{C}_{f_x}$. We actually show that $w \in \text{interior}(\mathcal{C}_{f_x})$.

Indeed, if this were not true, then

$$|Df_{f_x}^{-1}(w)| \geq m'_{\mathcal{C},x} |w|.$$

This would give

$$\begin{aligned}
|v| &= |Df_{f_x}^{-1}(w)| \geq m'_{C,x} |w| \\
&\geq m'_{C,x} \cdot m_{C,x} |v| \\
&> |v|
\end{aligned}$$

which is a contradiction. QED.

We say that f is *strongly dominating* on C over Λ if C has constant orbit core dimension and

$$(10) \quad \left(\inf_{x \in \Lambda} m_{C,x} \right) \cdot \left(\inf_{x \in \Lambda} m'_{C,x} \right) > 1.$$

Theorem 1.2. *Suppose that f is dominating on C over Λ . Then there is a unique Df -invariant splitting $T_\Lambda M = E_1 \oplus E_2$ such that for all $x \in \Lambda$, we have $E_{1x} \subset C_x$ and $E_{2x} \subset T_x M \setminus C_x$.*

Further, if f is strongly dominating on C over Λ , then the functions $x \rightarrow E_{1x}$, $x \rightarrow E_{2x}$ are continuous in x .

Proof.

Existence:

Let f be dominating on C .

Let $C_x^c = T_x M \setminus C_x$.

Set

$$C_x^+ = \bigcap_{n \geq 0} Df_{f^{-n}x}^n(C_{f^{-n}x}) \subset C_x,$$

$$C_x^- = \bigcap_{n \geq 0} Df_{f^n x}^{-n}(C_{f^n x}^c) = \bigcap_{n \geq 0} Df_{f^n x}^{-n}(\text{Closure}(C_{f^n x}^c)).$$

Let $E_{f^{-n}x}$ be the core of $C_{f^{-n}x}$.

Then, $E'_{n,x} = Df_{f^{-n}x}^n(E_{f^{-n}x})$ is a linear subspace which is contained in C_x , and they all have the same dimension.

Let

$$E_x = \lim_{n_i \rightarrow \infty} E'_{n_i,x}$$

in the Grassmann sense for some sequence $n_i \rightarrow \infty$.

Similarly, let $F_{f^n x}$ be a subspace contained in $C_{f^n x}^c$ complementary to the core of $C_{f^n x}$, and let $F'_{n,x} = Df_{f^n x}^{-n}(F_{f^n x})$.

We may assume that the sequence n_i is chosen such that $F_x = \lim_{n_i \rightarrow \infty} F'_{n_i,x}$.

Then, $E_x \subset C_x$, $F_x \subset \text{interior}(C_x^c)$, and $\dim E_x + \dim F_x = \dim T_x M$. Also, since $E_x \cap F_x = \{0\}$, we have $T_x M = E_x \oplus F_x$.

We claim:

$$(11) \quad C_x^- = F_x,$$

and

$$(12) \quad C_x^+ = E_x.$$

For $x \in \Lambda$, let $x_i = f^i x$, and set

$$m_{x_i} = m_{C_{\cdot, x_i}}(f), \quad m'_{x_i} = m'_{C_{\cdot, x_i}}(f).$$

For $n \geq 1$ set

$$M_n = \prod_{j=0}^{n-1} m_{x_j}, \quad M'_n = \left(\prod_{j=0}^{n-1} m'_{x_j} \right)^{-1}.$$

From condition (8) we may choose $\rho > 1$ so that

$$(13) \quad m_{x_j} \cdot m'_{x_j} > \rho \quad \forall j, \quad \text{which of course gives } \frac{M_n}{M'_n} > \rho^n.$$

Observe that for $v \in C_x^- \setminus \{0\}$, we have

$$(14) \quad Df_x^n(v) \in C_{x_n}^c \setminus \{0\},$$

and

$$(15) \quad |Df_x^n(v)| \leq M'_n |v|,$$

while, for $v \in C_x^+ \setminus \{0\}$, we have

$$(16) \quad Df_x^n(v) \in C_{f^n x} \setminus \{0\},$$

and

$$(17) \quad |Df_x^n(v)| \geq M_n |v|.$$

Let us now prove (11) and (12).

We know that $F_x \subset C_x^-$ and $E_x \subset C_x^+$. Suppose there is a $v \in C_x^- \setminus F_x$.

Write $v = v_1 + v_2$ with $v_1 \in E_x$, $v_2 \in F_x$. Of course, $v_1 \neq 0$.

From (15) and (17), we have

$$\begin{aligned} M'_n |v| &\geq |Df_x^n(v)| \\ &= |Df_x^n(v_1) + Df_x^n(v_2)| \\ &\geq M_n |v_1| - M'_n |v_2| \end{aligned}$$

or, by (13),

$$\frac{|v| + |v_2|}{|v_1|} \geq \frac{M_n}{M'_n} \geq \rho^n.$$

Since the left side of this equation is bounded, this is a contradiction, proving (11).

Replacing f by f^{-1} gives $C_x^+ = E_x$, so (12) is also proved.

Now, we set $E_{1x} = E_x$, $E_{2x} = F_x$ to give the splitting required in Theorem 1.2. It is clearly Df -invariant since both C_x^+ and C_x^- are.

Uniqueness:

If $\bar{E}_{1x} \subset C_x, \bar{E}_{2x} \subset C_x^c$ for all $x \in \Lambda$, and the \bar{E}_{ix} are Df -invariant, then we clearly have $\bar{E}_{1x} \subset C_x^+$ and $\bar{E}_{2x} \subset C_x^-$. Hence, $\bar{E}_{1x} \subset E_x$ and $\bar{E}_{2x} \subset F_x$. If $\dim \bar{E}_{1x} < \dim E_x$, then $\dim \bar{E}_{1x} \oplus \bar{E}_{2x} < \dim M$, contradicting the assumption that we have a splitting of $T_x M$. Hence, $E_x = \bar{E}_{1x}$. Similarly, $F_x = \bar{E}_{2x}$.

Continuity:

Assume that f is strongly dominating over C , and let $E \oplus F = \{E_x \oplus F_x\}$ be the induced Df -invariant splitting over Λ with $E_x \subset C_x, F_x \subset C_x^c$.

Suppose that $z \rightarrow F_z$ is not continuous at some x . We may choose a sequence y_i converging to x and subspaces \tilde{F}_x, \tilde{E}_x of $T_x M$ such that

$$(18) \quad \dim F_{y_i} \text{ is constant,}$$

$$(19) \quad F_{y_i} \rightarrow \tilde{F}_x \text{ and } E_{y_i} \rightarrow \tilde{E}_x \text{ in the Grassmann sense,}$$

and

$$(20) \quad \tilde{F}_x \neq F_x.$$

Case 1: There is a unit vector $v \in \tilde{F}_x \setminus F_x$.

Choose unit vectors $v_i \in F_{y_i}$ such that $v_i \rightarrow v$ as $i \rightarrow \infty$, and write $v = v_1 + v_2$ and $v_1 \in E_x, v_2 \in F_x$. Since v is not in F_x we have $v_1 \neq 0$.

For $n \geq 0$, let

$$M_n = \prod_{j=0}^{n-1} m_{f^j x}, \quad M'_n = \prod_{j=0}^{n-1} m'_{f^j x}, \quad M'_{n,i} = \prod_{j=0}^{n-1} m'_{f^j y_i}.$$

Now, for each $n > 0$, we have

$$\begin{aligned} |Df_x^n(v)| &= \lim_i |Df_{y_i}^n(v_i)| \\ &\leq \liminf_{i \rightarrow \infty} \frac{|v_i|}{M'_{n,i}} \\ &= \liminf_{i \rightarrow \infty} \frac{1}{M'_{n,i}}, \end{aligned}$$

and

$$\begin{aligned} |Df_x^n(v)| &= |Df_x^n(v_1) + Df_x^n(v_2)| \\ &\geq |Df_x^n(v_1)| - |Df_x^n(v_2)| \\ &\geq M_n |v_1| - \frac{1}{|M'_n|} |v_2|. \end{aligned}$$

Letting

$$m_0 = \inf_{x \in \Lambda} m_{C,x}(f), \quad m'_0 = \inf_{x \in \Lambda} m'_{C,x}(f),$$

we have

$$M'_{n,i} \geq (m'_0)^n, M_n \geq m_0^n, \text{ and } M'_n \geq (m'_0)^n.$$

Then, the above inequalities give

$$\liminf_{i \rightarrow \infty} \frac{1}{M'_{n,i}} \geq M_n |v_1| - \frac{|v_2|}{M'_n},$$

or,

$$\frac{1}{(m'_0)^n} + \frac{|v_2|}{(m'_0)^n} \geq m_0^n |v_1|,$$

or,

$$\frac{1 + |v_2|}{|v_1|} \geq (m'_0 m_0)^n.$$

For n large this is a contradiction.

Case 2: There is a unit vector $v \in F_x \setminus \tilde{F}_x$.

Write the unit vector v as $v = v_{1i} + v_{2i}$ with $v_{1i} \in E_{y_{n_i}}$ and $v_{2i} \in F_{y_{n_i}}$. The angles between the subspaces E_{y_i} and F_{y_i} are bounded below, so, for large i , the norms $|v_{1i}|$ are bounded below. Hence, the quantities

$$\frac{1 + |v_{2i}|}{|v_{1i}|}$$

are bounded above. On the other hand, as in Case 1, we can show that they are no smaller than $(m'_0 m_0)^n$ for each n . This contradiction proves that $z \rightarrow F_z$ is continuous.

Similarly, we obtain that $z \rightarrow E_x$ is also continuous.

This proves Theorem 1.2. QED.

Proposition 1.3. *A sufficient condition for f to have a dominated splitting over Λ is that there is an integer $n_0 > 0$ such that f^{n_0} has a strongly dominated cone field C over Λ .*

Proof.

By Theorem 1.2, we get a continuous Df^{n_0} -invariant splitting $T_x M = E_1 \oplus E_2$ and a $\tau > 1$ such that, for all $x \in \Lambda$,

$$m(Df_x^{n_0} | E_{1x}) \cdot m(Df_{f^{n_0}x}^{-n_0} | E_{2,f^{n_0}x}) > \tau.$$

This implies that for each $k \geq 0$,

$$m(Df_x^{kn_0} | E_{1x}) \cdot m(Df_{f^{kn_0}x}^{-kn_0} | E_{2,f^{kn_0}x}) > \tau^k.$$

Let $n \geq 0$ and write $n = kn_0 + r$ with $0 \leq r < n_0$.

Then,

$$(21) \quad m(Df_x^n | E_{1x}) \geq m(Df_x^{kn_0} | E_{1x}) \cdot \inf_{0 \leq j < n_0} m(Df_{f^{kn_0}x}^j | E_{1,f^{kn_0}x})$$

and

$$(22) \quad m(Df_{f^n x}^{-n} | E_{2,f^n x}) \geq m(Df_{f^{kn_0}x}^{-kn_0} | E_{2,f^{kn_0}x}) \cdot \inf_{0 \leq j < n_0} m(Df_{f^j x}^{-j} | E_{2,f^j x}),$$

so, we get

$$m(Df_x^n | E_{1x}) \cdot m(Df_{f^n x}^{-n} | E_{2, f^n x}) \geq C\tau^k \geq C_1\lambda^{n_0 k+r} = C_1\lambda^n$$

for some constants $C, C_1 > 0$ and $\lambda > 1$. This proves the Proposition. QED.

Let us now apply these results to give a useful condition for uniform hyperbolicity involving cone fields.

One standard definition of uniform hyperbolicity is the following.

Let $f : M \rightarrow M$ be a C^1 diffeomorphism of a Riemannian manifold M , and let Λ be a compact invariant subset of M ; i.e., $f(\Lambda) = \Lambda$. We say that Λ is a *uniformly hyperbolic* set for f , if there are constants $C > 0, \lambda > 1$ and a continuous splitting $T_x M = E_x^u \oplus E_x^s$ for each $x \in \Lambda$ such that

- (1) $Df_x(E_x^s) = E_{f_x}^s$ and $Df_x(E_x^u) = E_{f_x}^u$,
- (2) for $v \in E_x^u$ and $n \geq 0$, we have

$$|Df_x^n(v)| \geq C\lambda^n |v|,$$

- (3) for $v \in E_x^s$ and $n \geq 0$, we have

$$|Df_x^{-n}(v)| \geq C\lambda^n |v|.$$

Let us say that f is *expanding* on the cone field C if

$$\inf_{x \in \Lambda} m_{C,x}(f) > 1.$$

Similarly, we say that f is *co-expanding* on C if

$$\inf_{x \in \Lambda} m'_{C,x}(f) > 1.$$

We now use our previous results and definitions to get

Theorem 1.4. *A necessary and sufficient condition for Λ to be a uniformly hyperbolic set for f is that there are an integer $N > 0$ and a cone field C with constant orbit core dimension over Λ such that f^N is both expanding and co-expanding on C .*

Proof.

Sufficiency:

If C and N are as in the statement of the theorem, then, C is obviously strongly dominating for f^N over Λ . From Theorem 1.2 and Proposition 1.3 we get a dominated splitting $T_x M = E_x \oplus F_x$ for f over Λ with $E_x \subset C_x$ and $F_x \subset C_x^c$ for every $x \in \Lambda$. Now, using the Euclidean algorithm as in estimates (21) and (22), we get that $E_x \oplus F_x$ is a uniformly hyperbolic splitting for f on Λ .

Necessity:

Let Λ be uniformly hyperbolic with splitting $T_x = E_x^u \oplus E_x^s$ for $x \in \Lambda$. It is known [3] that there is an adapted Finsler norm $|\cdot|_1$ on TM . This is a norm induced by a Finsler metric such that there is a $\tau > 1$ such that for $x \in \Lambda$,

- (1) if $v \in E_x^u$, then $|Df_x(v)|_1 \geq \tau |v|_1$, and
- (2) if $v \in E_x^s$, then $|Df_x^{-1}(v)|_1 \geq \tau |v|_1$,
- (3) if $v = v_1 + v_2$ with $v_1 \in E_x^u, v_2 \in E_x^s$, then $|v|_1 = \max(|v_1|_1, |v_2|_1)$.

For $x \in \Lambda$, set

$$C_x = \{v = (v_1, v_2) \in E_x^u \oplus E_x^s : |v_1|_1 \leq |v_2|_1\}.$$

The core of C_x may be taken to be E_x^u . This is clearly an expanding and co-expanding cone field of constant core dimension for f relative to the adapted norm $|\cdot|_1$. But, since any two norms have uniformly bounded ratios on non-zero vectors, it is clear that in any norm there is a positive integer N such that C is expanding and co-expanding for f^N . QED.

2. APPLICATION TO THE HENON FAMILY

In this section we apply the preceding results to two dimensional systems, both real and complex. This gives a simple proof of hyperbolicity on the set of bounded orbits in the Henon family $H(x,y) = (A - x^2 - By, x)$ for $|A|$ sufficiently large depending on $|B|$. In the real case this was originally proved by Devaney and Nitecki [2]. Their proof is elementary but heavily dependent on real geometry. In fact, it is not clear that their methods apply to the complex case. Before we began the work reported here, we asked several experts about the complex case and were told that R. Oberste-Vorth has a proof which, however, was never published. We do not know what estimates he obtained for the relative sizes of $|A|$ and $|B|$.

Devaney and Nitecki proved that hyperbolicity holds provided that

$$(23) \quad A > \left(\frac{5 + 2\sqrt{5}}{4} \right) (1 + |B|)^2.$$

At the end of our calculations below, we were, in fact, quite surprised to find that essentially the same estimate holds in the complex case. In retrospect, it seems that our arguments are very similar to those in [2], so our main contribution here, perhaps, is that we have made the arguments cleaner and removed the dependence on real geometry. After this work was done, John Smillie informed us that better estimates can be obtained using complex methods. In fact, Proposition 7.4.6 in [5] gives that the bounded orbits form a topological horseshoe for $A > 2(1 + |B|)^2$, and then complex methods can be used to show actual hyperbolicity. Nevertheless, our estimates are sufficiently elementary that we feel it is desirable to present them here.

As is well-known, all degree two polynomial diffeomorphisms are affinely conjugate to members of the Henon family. Hence, in studying their dynamical properties, there is no loss of generality in assuming that $0 < |B| \leq 1$

Theorem 2.1. *Consider the real or complex Henon family $H_{A,B}(x,y)$ with $0 < |B| \leq 1$. Let $\Lambda_{A,B}$ denote the set of points with bounded orbits. Assume that*

$$(24) \quad |A| > \left(\frac{5 + 2\sqrt{5}}{4} \right) (1 + |B|)^2.$$

In the complex case or the real case with $A > 0$, we have that $\Lambda_{A,B}$ is a non-empty compact invariant uniformly hyperbolic set. In the real case with $A < 0$, the set $\Lambda_{A,B}$ is empty.

Let us proceed to the proof.

Consider $V = \mathbf{R}$ or $V = \mathbf{C}$ and the family of maps $H(x,y) = H_{A,B}(x,y)$ for $x,y \in V$. If $V = \mathbf{R}$, we let A, B be real, but we allow them to be complex if $V = \mathbf{C}$.

We begin with simple criteria for expansion and co-expansion of a linear map using the standard unit cone. We use the max norm

$$|v| = \max(|v_1|, |v_2|)$$

for a vector $v = (v_1, v_2) \in V^2$.

Define the standard unit cone, \mathcal{K} in $V^2 = V \oplus V$.

$$\mathcal{K} = \{v = (v_1, v_2) : |v_2| \leq |v_1|\}.$$

The complementary cone is

$$\mathcal{K}^c = \{v = (v_1, v_2) : |v_1| < |v_2|\}.$$

For a linear map $L : V^2 \rightarrow V^2$ with matrix

$$L = \begin{pmatrix} L_{11} & L_{12} \\ L_{21} & L_{22} \end{pmatrix}$$

and determinant $\det(L) = L_{11}L_{22} - L_{12}L_{21}$, we have the inverse matrix

$$L^{-1} = \frac{1}{\det(L)} \begin{pmatrix} L_{22} & -L_{12} \\ -L_{21} & L_{11} \end{pmatrix}.$$

Lemma 2.2. *A sufficient condition for L to be expanding on \mathcal{K} is that*

$$(25) \quad |L_{11}| - |L_{12}| > 1.$$

Analogously, a sufficient condition for L to be co-expanding on \mathcal{K} is that

$$(26) \quad |L_{11}| - |L_{21}| > |\det(L)|.$$

Proof.

Let $\lambda > 1$ be such that

$$|L_{11}| - |L_{12}| > \lambda.$$

For $v = (v_1, v_2) \in \mathcal{K}$, we have $|v| = |v_1|$, and

$$\begin{aligned} |Lv| &\geq |L_{11}v_1 + L_{12}v_2| \\ &\geq |L_{11}| |v_1| - |L_{12}| |v_2| \\ &= |v_1| (|L_{11}| - |L_{12}| \frac{|v_2|}{|v_1|}) \\ &\geq \lambda |v_1| \\ &= \lambda |v|. \end{aligned}$$

Similarly, let $\lambda > 1$ be such that

$$|L_{11}| - |L_{21}| > \lambda |\det(L)|.$$

For $v \in \mathcal{K}^c$, we have $|v| = |v_2|$, and

$$\begin{aligned}
 |L^{-1}v| &\geq \frac{1}{|\det(L)|} (|-L_{21}v_1 + L_{11}v_2|) \\
 &\geq \frac{1}{|\det(L)|} (|L_{11}v_2| - |L_{21}v_1|) \\
 &\geq \frac{1}{|\det(L)|} |v_2| (|L_{11}| - |L_{21}| \frac{|v_1|}{|v_2|}) \\
 &\geq \lambda |v_2| \\
 &= \lambda |v|.
 \end{aligned}$$

QED.

Now, we return to the Henon family $H = H_{A,B}(x,y) = (A - x^2 - By, x)$ with $0 < |B| \leq 1$.

For $C > 0$ let $Q_C = \{(x,y) \in V^2 : \max(|x|, |y|) \leq C\}$.

In the real case, the next result is contained in [2].

Proposition 2.3. *Let C_0 be the largest root of the quadratic polynomial*

$$p(r) = r^2 - (1 + |B|)r - |A|,$$

and, let $Q_0 = Q_{C_0}$. Then,

$$\Lambda_{A,B} \subset Q_0.$$

Proof.

Write $(x_1, y_1) = (A - x^2 - By, x)$.

Take any $C > C_0$. We show that $\Lambda_{A,B} \subset Q_C$. The result then follows taking the intersection

$$\bigcap_{C > C_0} Q_C.$$

Since C is greater than the largest root of the polynomial

$$r^2 - (1 + |B|)r - |A|,$$

we can choose $\lambda > 1$ close enough to 1 so that C is greater than the largest roots of the polynomials

$$p_1(r) = r^2 - (\lambda + |B|)r - |A|,$$

and

$$p_2(r) = r^2 - (1 + \lambda|B|)r - |A|.$$

Consider the sets $\mathcal{K}^+ = \mathcal{K} \setminus Q_C$, and $\mathcal{K}^- = \mathcal{K}^c \setminus Q_C$.

Let $\pi_1(x,y) = x$, $\pi_2(x,y) = y$ be the natural projections on \mathbf{R}^2 , and write $H = H_{A,B}$.

We show

- (a) $H(\mathcal{K}^+) \subset \mathcal{K}^+$ and $z \in \mathcal{K}^+$ implies that $|\pi_1 H(z)| > \lambda |z|$, and
- (b) $H^{-1}(\mathcal{K}^-) \subseteq \mathcal{K}^-$ and $z \in \mathcal{K}^-$ implies that $|\pi_2 H^{-1}(z)| > \lambda |z|$.

These statements give that the norms of the forward iterates of points in \mathcal{K}^+ approach infinity while the norms of the backward iterates of points in \mathcal{K}^- approach infinity. Hence, bounded orbits must be in Q_C .

If $(x,y) \in \mathcal{K}^+$, then $|y| \leq |x|$ and $|x| > C$, so

$$\begin{aligned}
|x_1| &= |A - x^2 - By| \\
&\geq |x|^2 - |A| - |B||y| \\
&\geq |x|^2 - |A| - |B||x|
\end{aligned}$$

For $|x_1| > \lambda|x|$, we need

$$|x|^2 - |A| - |B||x| > \lambda|x|,$$

or, $|x|$ must be greater than the largest root of $p_1(r)$. But, this follows from our choice of λ since $|x| > C$.

Further, since, $|y_1| = |x| < |x_1|$, we have $H(x, y) \in \mathcal{K}^+$.

This is statement (a).

Statement (b) is similar.

Suppose $(x_1, y_1) \in \mathcal{K}^-$.

Then, $|x_1| < |y_1|$ and $|y_1| > C$ and

$$(x, y) = H^{-1}(x_1, y_1) = \left(y_1, \frac{A - y_1^2 - x_1}{B}\right)$$

give

$$\begin{aligned}
|y| &= \frac{1}{|B|} |A - y_1^2 - x_1| \\
&\geq \frac{1}{|B|} (|y_1|^2 - |x_1| - |A|) \\
&\geq \frac{1}{|B|} (|y_1|^2 - |y_1| - |A|).
\end{aligned}$$

To get $|y| > \lambda|y_1|$ we need that $|y_1|$ be larger than the largest root of $p_2(r)$.

Again, this follows from the choice of λ since $|y_1| > C$. In addition, we have $|x| = |y_1| < |y|$, so $(x, y) \in \mathcal{K}^-$. QED.

It remains to prove the hyperbolicity of $\Lambda_{A,B}$ under the assumption (24).

Let

$$\alpha = 1 + |B|,$$

and, let

$$C_0 = \frac{1}{2} \left(\alpha + \sqrt{\alpha^2 + 4|A|} \right)$$

be as in Proposition 2.3.

An elementary algebraic calculation shows that the inequality in (24) is equivalent to that in (27). So, Theorem 2.1 follows from

Proposition 2.4. *In the complex Henon map, suppose*

$$(27) \quad |A| > \frac{\alpha^2}{4} + \alpha C_0,$$

Then, $\Lambda_{A,B}$ is a non-empty hyperbolic set for $H_{A,B}$.

In the real Henon map, if (27) holds, then $\Lambda_{A,B}$ is hyperbolic if $A > 0$ and empty if $A < 0$.

Proof.

For notational ease, let us suppress the subscripts and write $H = H_{A,B}$, and $\Lambda = \Lambda_{A,B}$.

We will prove that

(\star) *If (27) holds, then Λ is hyperbolic provided that it is not empty.*

This takes care of the complex case. (In that case, $\Lambda_{A,B}$ is clearly non-empty since H has periodic points.)

For the real case, let U be the set of parameters $(A, B) \in \mathbf{R}^2$ satisfying (24), and let $G \subset U$ be the subset of those $(A, B) \in U$ such that $\Lambda_{A,B} \neq \emptyset$.

Letting

$$\beta = \left(\frac{5 + 2\sqrt{5}}{4} \right) \alpha^2$$

be the number on the right side of the inequality (24), we have that the set U is the disjoint union of the two open connected sets

$$U_1 = \{(A, B) : A > \beta\},$$

and

$$U_2 = \{(A, B) : -A < -\beta\}.$$

Statement (\star) and the local stability of hyperbolic isolated invariant sets (Theorem 3, page 157 in [8]) imply that G is an open subset of U . We claim $U \setminus G$ is also open.

Indeed, we have

$$\Lambda = \bigcap_n H^n(Q_0)$$

is the decreasing intersection of the compact sets

$$\bigcap_{-k \leq n \leq k} H^n(Q_0).$$

If Λ were empty, then for some finite $k_0 > 0$, we would have

$$\bigcap_{-k_0 \leq n \leq k_0} H^n(Q_0) = \emptyset,$$

and this property persists under small changes in (A, B) .

It follows that $G \cap U_i$ is either empty or equal to U_i for $i = 1, 2$.

Fix $0 \leq |B| \leq 1$. Then,

$$|C_0| \leq 1 + \sqrt{1 + |A|}$$

Using this, it is easy to show that if A is very large and negative, then for any $(x, y) \in Q_0$, we have

$$\pi_1 H^n(x, y) \rightarrow -\infty$$

as $n \rightarrow \infty$. Thus, if $|A|$ very large and negative, we have $\Lambda = \emptyset$. Hence, Λ must be empty for all A in $(-\infty, -\beta)$. On the other hand, if A is very large and positive, then Λ is non-empty since $H_{A,B}$ has fixed points. So, Λ is non-empty for all A in (β, ∞) .

It remains to prove (\star) .

Now,

$$DH = \begin{pmatrix} -2x & -B \\ 1 & 0 \end{pmatrix} \text{ and } \det(DH) = B,$$

so, (25) and (26) are both implied by

$$|2x| > 1 + |B|,$$

or,

$$(28) \quad |x| > \frac{1 + |B|}{2} = \frac{\alpha}{2}.$$

We want to consider $z = (x, y) \in Q_0$ such that

$$(29) \quad |x| \leq \frac{\alpha}{2}.$$

We show that (29) together with (27) implies that the image $H(z)$ is not in Q_0 . This will imply that \mathcal{X} is expanded and co-expanded by DH on $Q_0 \cap H^{-1}(Q_0)$. Since Λ is contained in this latter set, this will prove hyperbolicity.

To get $H(z) \notin Q_0$, it suffices to have

$$|A - x^2 - By| > C_0.$$

Since $|x| \leq \frac{\alpha}{2}$ and $|y| \leq C_0$, this is implied by

$$|A| - \frac{\alpha^2}{4} - |B|C_0 > C_0,$$

or

$$(30) \quad |A| - \frac{\alpha^2}{4} > \alpha C_0.$$

But this is the same as (27). QED.

Remark. In addition to the hyperbolicity, Devaney and Nitecki prove that, in the real case with $\frac{A}{|B|}$ large, the set of bounded orbits of $H_{A,B}$ equals its non-wandering set, and the map $H_{A,B}$ restricted to this set is topologically conjugate to the full two-sided shift on two symbols. We easily obtain this in our situation even in the complex case. First note that these facts are fairly easy to prove for a real Henon map H_{A_1, B_1} with A_1 large and $|B_1|$ sufficiently small. In the general (even complex) case, for arbitrary (A, B) satisfying (27), we connect $H_{A,B}$ to a real Henon map H_{A_1, B_1} through a curve $\{H_t = H_{A_t, B_t}, t \in [0, 1]\}$ of Henon maps keeping (27) and $\Lambda_t \neq \emptyset$ for all t . Then, each Λ_t is a hyperbolic isolated invariant set. Hence, the local stability of isolated invariant hyperbolic sets gives that all the pairs (H_t, Λ_t) are topologically conjugate.

REFERENCES

- [1] E. Bedford and J. Smillie. Real Polynomial Diffeomorphisms with Maximal Entropy: Tangencies. <http://front.math.ucdavis.edu/math.DS/0103038>, 1, 6 March, 2001.
- [2] R. Devaney and Z. Nitecki. Shift automorphism in the henon mapping. *Commun. Math. Phys.*, 67:137–148, 1979.
- [3] M. Hirsch and C. Pugh. Stable manifolds and hyperbolic sets. *Proc. AMS Symp. Pure Math.*, 14, 1970.
- [4] R. Mane. A proof of the C^1 stability conjecture. *Publ. Math. I.H.E.S.*, 66:161–210, 1988.
- [5] S. Morosawa, Y. Nishimura, M. Taniguchi, and T. Ueda. *Holomorphic Dynamics*. Cambridge University Press, 2000.
- [6] J. Moser. *Stable and Random Motions in Dynamical Systems*. Annals of Math. Studies. Princeton University Press, 1973.
- [7] S. Newhouse and J. Palis. Bifurcations of Morse-Smale dynamical systems. In M. M. Peixoto, editor, *Dynamical Systems: Proc. Symp. Bahia, Brazil, July 26-Aug. 14, 1971*, pages 303–366. Academic Press, 1973.
- [8] J. Palis and F. Takens. *Hyperbolicity and Sensitive Chaotic Dynamics at Homoclinic Bifurcations.*, volume 35 of *Cambridge Studies in Advanced Mathematics*. Cambridge University Press, 1993.
- [9] E. Pujals and M. Sambarino. Homoclinic tangencies and hyperbolicity for surface diffeomorphisms. *Annals of Math.*, 151:962–1023, 2000.
- [10] Clark Robinson. *Dynamical Systems, Stability, Symbolic Dynamics, and Chaos, second ed.* Studies in Advanced Mathematics. CRC Press, 1999.